3.8: Rewrite Equations and Formulas

Goals: *Isolate a variable in a literal equation

*Rewrite an equation so it is in function form

Literal equation: an equation in which coefficients and constants have been replaced by letters


Ex: Solve ax + b = c for x This means to: isolate x

Solve the following equations for the given variable.

Ex: Solve $A = \frac{1}{2}bh$ for h

$$2A = bh$$
 To get rid of $\frac{1}{2}$ multiply by 2 $\frac{2A}{b} = h$

Find h if the shown triangle has an area of 64.4 m²



Plug in A and b into new equation. Plug numbers to solve for h

$$\frac{2(64.4)}{14} = h$$

$$\frac{128.8}{14} = h$$

$$9.2 \text{ m} = h$$

Ex: p + qx = r for x

$$\frac{r-p}{a} = x$$

Ex: A = lw for l, then find l if A = 351 cm² and w = 13 cm

$$\frac{A}{w} = l$$

$$\frac{351}{13} = l$$
 $l = 27 \text{ cm}$

Ex: You are visiting Toronto over the weekend and look up a weather forecast. Find the low temperatures for Saturday and Sunday in degrees Fahrenheit. First rewrite the conversion formula so *F* is isolated:

$$C = \frac{5}{9}(F - 32)$$

	Friday	Saturday	Sunday
Forecast	Sunny	Sunny	Partly Cloudy
High	21° <i>C</i>	22°C	16° <i>C</i>
Low	13° <i>C</i>	14° <i>C</i>	10° <i>C</i>

$$\frac{9}{5}C+32=F$$

Saturday

$$\frac{9}{5} \cdot 14 + 32 = F$$

$$57.2 = F$$

Sunday

$$\frac{9}{5} \cdot 10 + 32 = F$$

$$50 = F$$

**RECALL THAT ALL FUNCTIONS START WITH: y = **

So when you are rewriting an equation so it is in function form that means to isolate: y =

Ex: -2x + 3y = 6

$$\frac{3y}{3} = \frac{6+2x}{3}$$

$$y = 2 + \frac{2}{3}x$$

Ex: 3x + 2y = 8

$$\underline{2y} = \underline{8 - 3x}$$

$$y = 4 - \frac{3}{2}x$$

Ex: 4x - 2y = -6

$$\frac{-2y}{-2} = \frac{-6 - 4x}{-2}$$

$$y = 3 + 2x$$

Ex: -3x - y = 7

$$\frac{-y}{-1} = \frac{7+3x}{-1}$$

$$y = -7 - 3x$$