

# Ch 18 Surgical Asepsis Test Guidelines

## 15 minute Skill

- NO Jewelry/watch during test
- Verbalize that bedside table has been disinfected
- Verbalization of Surgical Scrub
- Supplies: Foley Cath Kit (sterile field), Sterile 4 X4 Gauze, Sterile Solution, Sterile Container (Specimen cup), Sterile Gloves
- Table at waist level or higher
- Check ALL supplies for expiration dates, no moisture, penetration or breakage of items
- Keep items 1 inch from the edges. Vertical sides are NOT sterile, Horizontal are.
- Open Foley Kit-Distal, Lateral, Lateral, Proximal.
- Open Sterile Container-Keeping lid sterile at all times
- Open Sterile solution with Label in palm of the hand. Pour solution into sterile cup NOT contaminating Sterile Lid and no splashing of solution
- Open gauze distal to proximal using flaps making sure flaps DO NOT fold back
- Open Sterile gloves, using the bottom corner, open top folds then bottom folds, grab center fold to expose gloves
- Don Sterile gloves without contamination
- Pick up gauze with sterile gloves
- Fold one gauze and dip in sterile solution, ring out sterile gauze
- Place wet gauze on sterile Foley kit
- Place dry gauze on sterile drape
- Remove sterile gloves without contamination

