

WORK SHEET

DIRECTIONS: The medication order is listed at the beginning of each problem. Calculate the oral doses. Show your work. Shade each medicine cup or oral syringe when provided to indicate the correct dose.

1. The licensed prescriber orders Minipress 2 mg po two times a day for Mr. Shaw's high blood pressure. How many capsules will the nurse administer per dose? _____

NDC0663-4310-71

250 Capsules

Minipress®
prazosin hydrochloride

1 mg[†]

CAUTION: Federal law prohibits dispensing without prescription.

 Manufactured by
Pfiizer Pharmaceuticals, Inc., Bartlesville, OK 74007

LABORATORIES DIVISION
New York, N.Y. 10017

RECOMMENDED STORAGE
STORE BELOW 85° F (30° C)
Dispense in light-resistant
containers (USP).
† Each capsule contains prazosin
hydrochloride equivalent to
1 mg of prazosin.



READ ACCOMPANYING
PROFESSIONAL INFORMATION
DOSAGE: See accompanying
prescribing information.
IMPORTANT: This closure is
not child-resistant.

6505-01-039-6320

4444

2. Mrs. Taylor has a long history of seizures. Elixir of phenobarbital 30 mg po q 12 h is ordered. How many milliliters will the nurse administer per dose? _____

NDC 0002-2438-05
16 fl oz (1 pt) (473 ml)

**ELIXIR No. 227
PHENOBARBITAL
ELIXIR, USP**

Contains 20 mg Phenobarbital
per 5 ml

WARNING—May be habit forming.
Contains Alcohol 14 Percent

CAUTION—Federal (U.S.A.) law
prohibits dispensing without
prescription.

Keep Tightly Closed
Store at Controlled Room Temperature 59° to 86° F
(15° to 30° C)
NPL 5991 AAIX
Eli Lilly & Company, Indianapolis, IN 46285, U.S.A.
Expiration Date/Control No.

VOID

Usual Adult Sedative Dose—60 mg (1 gr) (3 tea-
spoonfuls) two to four times a day.
Indiscriminate use may be harmful.
Dispense in a light-resistant container.


3 0002 2438 05 9

3. The licensed prescriber orders Crystodigin 0.2 mg po two times a day for 4 days, then 0.15 mg po two times a day. You have Crystodigin 0.05-mg tablets available. How many tablets will you give for each dose the first 4 days? _____ How many tablets will you give for each dose thereafter? _____

4. Mr. Davis has a diagnosis of acute maxillary sinusitis. His licensed prescriber orders Biaxin 500 mg q 12 h \times 10 days. How many tablets will the nurse administer per dose? _____

Exp. Lot	03-2185-3/R5	©Abbott	NDC 0074-8368-60 60 Tablets	6505-01-354-8582 Do not accept if seal over bottle opening is broken or missing. Dispense in a USP tight, light-resistant container. Each tablet contains: 250 mg clarithromycin. Usual Adult Dose: One or two tablets every twelve hours. See enclosure for full prescribing information. FilmTab - Film-sealed tablets, Abbott, Abbott Laboratories North Chicago, IL 60064, U.S.A.
	007436860		BIAXIN® FILMTAB® clarithromycin tablets 250 mg	
	Store tablets at 15° to 30°C (59° to 86°F).	NEW TABLETS	Caution: Federal (U.S.A.) law prohibits dispensing without prescription.	

5. Mrs. Rios complains of nausea. Compazine 2.5 mg po three times a day is ordered. The stock supply is Compazine syrup 5 mg/5 mL. How many milliliters will the nurse administer per dose? _____



6. The licensed prescriber orders Ativan 2 mg at bedtime. How many tablets will the nurse administer per dose? _____


1001 VG 78000000 5-20-9001	Ativan® (lorazepam) 1 mg 100 tablets	NDC 0008-0064-02 6505-01-057-9046	LOT EXP
Keep tightly closed (protect from moisture). Protect from light. Do not store above 86°F (30°C). Bristol-Myers Squibb Company Princeton, NJ 08543 USA 515405DRL-2	SEAL FOR PROTECTION	Use: Orally. One to 2 mg per day, given in divided doses. For information on interactions, see package insert.	

7. The licensed prescriber orders Pravachol 20 mg po at bedtime. How many tablets will the nurse administer per dose? _____

Exp. Cont.:	Keep tightly closed (protect from moisture). Protect from light. Do not store above 86°F (30°C). NDC 0003-5154-05 Bristol-Myers Squibb Company Princeton, NJ 08543 USA 515405DRL-2	90 tablets	NDC 0003-5154-05	NEW SMALLER TABLET	Each tablet contains 10 mg pravastatin sodium. Dosage: See insert. US Patents: 4,346,227; 5,030,447; 5,180,589
		10 mg	PRAVACHOL® Pravastatin Sodium Tablets		
			Caution: Federal law prohibits dispensing without prescription		
			Bristol-Myers Squibb Company		

8. Mandelamine 1 g po four times a day is scheduled for Mr. Eaton to treat his urinary tract infection. You have 0.5-g tablets available. How many tablets will you administer per dose? _____

9. The licensed prescriber orders Prozac 40 mg po daily in AM. How many milliliters will the nurse administer per dose? _____



NDC 0777-5120-58
120 mL M-5120

PROZAC®
LIQUID
FLUOXETINE
HYDROCHLORIDE
ORAL SOLUTION


Equivalent to
20 mg per 5 mL
Base

See accompanying literature for dosage.
Dispense in a light, light-resistant container.
Each 5 mL contains Fluoxetine Hydrochloride
equivalent to 20 mg Fluoxetine base
Contains alcohol 0.23%

Keep Tightly Closed
Rx only

Store at Controlled Room Temperature 59° to 86°F
(15° to 30°C)
N6 650 DPX

Eli Lilly and Company
Indianapolis, IN 46285, USA
Expiration Date/Control No.



10. Mr. Chang has Parkinson's disease and is to receive Cogentin 1 mg po at 1900. How many tablets will the nurse administer per dose? _____



NDC 0006-0021-68


100 TABLETS
COGENTIN® 0.5 mg
(BENZTROPINE MESYLATE)

Merck & Co., Inc.
West Point, PA 19486, USA

See accompanying circular.
Dispense in a well-closed container.
CAUTION: Fodder (USA) containing benzotropine mesylate may be hazardous to the environment.

This is a bulk package and not intended for dispensing.

11. Mrs. Martin receives Motrin 800 mg po three times a day for arthritis pain. The drug is supplied in 400-mg tablets. How many tablets will the nurse administer per dose? _____



See package insert for complete product information.

Dispense in light, light-resistant container.

Store at controlled room temperature 15° to 30° C (59° to 86° F).

610 012 103

The Upjohn Company
Kalamazoo, MI
49001, USA

Motrin®
Ibuprofen
Tablets, USP
400mg

NDC 0025-0120-15
150 Tablets

See package insert for dosage and full prescribing information.


Dispense in a light container with child-resistant closure.

STORAGE: CONTROLLED ROOM TEMPERATURE

Upjohn Pharmacy, Inc.
Subsidiary of
Schein Pharmaceutical, Inc.
Fortham Park, NJ 07932, USA

12. The licensed prescriber orders lithium carbonate 0.6 g po two times a day. The drug is supplied in 300-mg scored tablets. How many tablets will the nurse administer per dose? _____ How many milligrams will be given each day? _____

13. Your patient is taking minoxidil 30 mg daily for hypertension. How many tablets will you administer per dose? _____



SCHEIN
PHARMACEUTICAL

NDC 0364-2173-01 180 Tablets

MINOXIDIL
Tablets, USP
10 mg

See package insert for dosage and full prescribing information.

Dispense in a light container with child-resistant closure.

STORAGE: CONTROLLED ROOM TEMPERATURE

Upjohn Pharmacy, Inc.
Subsidiary of
Schein Pharmaceutical, Inc.
Fortham Park, NJ 07932, USA

0364-2173-01
Control Number and Expiration Date

SAMPLE

14. Mr. Hill is to receive Cipro 0.75 g q 12 h for a knee infection. How many tablets will the nurse administer per dose? _____

051438 NDC 0179-1226-50

CIPRO®
(ciprofloxacin hydrochloride)
Equivalent to
750 mg ciprofloxacin
50 Tablets
Caution: Federal (USA) law prohibits dispensing without a prescription.

VOID

Bayer
Bayer Corporation
Pharmaceutical Division
490 Morgan Lane
West Haven, CT 06516

DESCRIPTION: Each tablet contains ciprofloxacin hydrochloride equivalent to 750 mg of ciprofloxacin.
DOSAGE: See accompanying literature for complete information on dosage and administration.
RECOMMENDED STORAGE: Store below 86°F (30°C).

Batch: **VD D**
Expires: **VD D**

Printed in USA
©1995 Bayer Corporation 5402
PL500037

15. The licensed prescriber orders Geodon 80 mg bid to treat Mrs. Basey's acute agitation. You have 40-mg capsules available. How many capsules will you administer per dose? _____

Store in controlled room temperature, 15° to 30°C (59° to 86°F).

60 Capsules NDC 0049-3970-60

Geodon™
(ziprasidone HCl)
40 mg*

Pfizer Roerig
Division of Pfizer Inc, NY, NY 10017

DOSAGE AND USE
See accompanying prescribing information.

*Each capsule contains ziprasidone hydrochloride monohydrate equivalent to 40 mg ziprasidone.

Rx only

MADE IN USA

16. The licensed prescriber orders Gantrisin 4 g po STAT, then 2 g q 6 h. How many tablets will be given for the STAT dose? _____ How many tablets will be given for each of the 2-g doses? _____

NDC 0004-0009-01 ITEM 73314

Roche

GANTRISIN®
(sulfisoxazole)
0.5 g

Each tablet contains 0.5 g sulfisoxazole.
CAUTION: Federal law prohibits dispensing without prescription.

100 Tablets

USUAL DOSAGE: For dosage recommendations and other important prescribing information, read accompanying insert.
ROCHE LABORATORIES
a division of Hoffmann-La Roche Inc., Nutley, NJ 07110

Dispense in tight, light-resistant containers as defined in USP/NF.
STORE AT 59° TO 86° F (15° TO 30° C).
LOT
EXPIRES

0293

17. The licensed prescriber orders acyclovir 800 mg po q 4 h while the patient is awake. Acyclovir 400-mg tablets are available. How many tablets will the nurse administer per dose? _____

18. The licensed prescriber orders Gaviscon 30 mL po four times a day. Gaviscon is supplied in 360-mL bottles. How many mLs will be given in 1 day? _____

19. Ms. Vega complains of a rash on her abdomen, and the licensed prescriber orders Benadryl 30 mg po three times a day. How many milliliters will the nurse administer per dose? _____

N 0071-2220-17

ELIXIR

Benadryl®
(Diphenhydramine Hydrochloride Elixir, USP)

Caution—Federal law prohibits dispensing without prescription.

4 FLUIDOUNCES

PARKE-DAVIS
Div of Warner-Lambert Co
Morris Plains, NJ 07950 USA

Elixir P-D 2220 for prescription dispensing only.

Contains—12.5 mg diphenhydramine hydrochloride in each 5 mL. Alcohol, 14%.

Dose—Adults, 2 to 4 teaspoonfuls; children over 20 lb, 1 to 2 teaspoonfuls; three or four times daily.

See package insert.

Keep this and all drugs out of the reach of children.

Store below 30°C (86°F). Protect from freezing and light.

Exp date and lot

2220G102

20. Mr. Gifford has had a lumbar laminectomy and requires pain medication. The patient has an order for codeine 60 mg po q 3 h prn. How many tablets will the nurse administer per dose? _____

100 No. 250

POISON

SOLUBLE TABLETS

CODEINE PHOSPHATE 30

SOLUBLE TABLETS

30 mg

Warning—May be habit forming.

For Prescription Compounding Use Only

Usual Adult Dose—30 mg every 4 hours, q.s.

Contains Lactose and Sucrose.

Dispense in a light-resistant container.

N 0002-2557-02

21. The licensed prescriber orders Amoxil suspension 5 5 mL (125 mg/5 mL strength) po q 6 h for your patient who had a tonsillectomy. How many milligrams will you administer every 6 hours? _____

AMOXIL®
125mg/5mL

125mg/5mL
NDC 0029-6008-23

AMOXIL®
AMOXICILLIN FOR ORAL SUSPENSION

100mL
(when reconstituted)

SB SmithKline Beecham

Directions for mixing: Tap bottle until all powder flows freely. Add approximately 1/3 total amount of water for reconstitution. (Total=70 mL) shake vigorously to wet powder. Add remaining water; again shake vigorously. Each 5 mL (1 teaspoonful) will contain amoxicillin trihydrate equivalent to 125 mg amoxicillin.

Usual Adult Dosage: 250 to 500 mg every 8 hours.

Usual Child Dosage: 20 to 40 mg/kg/day in divided doses every 8 hours, depending on age, weight and infection severity. See accompanying prescribing information.

Keep tightly closed. Shake well before using. Refrigeration preferable but not required. Discard suspension after 14 days.

LOT: 9405793-F

EXP:

137.5 mg

70 mL / 125 mg / 5 mL = 14 / 5 = 2.8


2.8 x 100 mL = 280 mL

280 mL / 2 = 140 mL


140 mL / 2 = 70 mL

25 mg

22. Mr. Sawyer is admitted with congestive heart failure. His orders require Lasix 80 mg po daily. How many tablets will the nurse administer per dose? _____



23. The licensed prescriber orders nitroglycerin 0.4 mg sublingual prn for angina. The patient should take no more than three tablets in 15 minutes. Mr. Cane took two tablets 5 minutes apart. How many milligrams did he receive? _____



24. Mr. Koehler has rheumatoid arthritis and has Decadron 1.5 mg po q 12 h ordered. You have Decadron elixir 0.5 mg/5 mL. How many milliliters will you administer per dose? _____
How many ounces will you administer per dose? _____

$$\begin{array}{r} 3 \\ \hline 1.5 \text{ mg} \times 5 \text{ mL} = 15 \text{ mL} \\ 0.5 \text{ mg} \end{array}$$

25. Your adult patient has acute bronchitis and has cefaclor 500 mg po q 12 h ordered. How many milliliters will you administer per dose? _____

NDC 0002-5132-48
100 mL (When Mixed) M-5132

CECLOR®
CEFACLOR
FOR ORAL
SUSPENSION, USP
375 mg
per 5 mL

CAUTION—Federal (USA) law prohibits dispensing without prescription.

Usual Dose—Pediatric patients, 20 mg per kg a day (40 mg per kg in colic media) in two divided doses. Adults, 375 mg two times a day. See literature for complete dosage information. Contains Cefaclor Monohydrate equivalent to 7.5 g anhydrous Cefaclor in a 50 mL plastic bottle. Store at room temperature (20° to 25°C) or in controlled room temperature (15° to 30°C). Directions for Mixing—Add 62 mL of water in two portions to the dry mixture in the bottle. Shake well after each addition. mL (Approx. one teaspoonful) will then contain Cefaclor Monohydrate equivalent to 375 mg anhydrous Cefaclor. VVY 4525 AMX Aided by Eli Lilly Industries, Inc., Corcoran, Puerto Rico 00918 a subsidiary of Eli Lilly & Co., Indianapolis, IN, USA. Expiration Date

100 mL CECLOR® CEFACLOR FOR ORAL SUSPENSION, USP 375 mg per 5 mL. Opened bottle provide extra space for shaking. Shake well before use. Do not use if the powder has become clumpy. Keep tightly closed. Discard unused portion after 14 days. SHAKE WELL BEFORE USING

Control No.

26. Mrs. Turner is admitted with hypertension. Apresoline 25 mg po four times a day is ordered. You have 50-mg scored tablets available. How many tablets will you administer per dose? _____

27. Your patient complains of indigestion during meals. Mylanta 30 mL po pc four times a day is ordered. Mylanta is supplied in a 360-mL bottle. There are _____ doses in one 360-mL bottle.

28. The licensed prescriber orders Halcion 0.25 mg po at bedtime. How many tablets will the nurse administer per dose? _____

Pharmacist: Dispense in this container with patient leaflet attached. See package insert for complete product information. Keep container tightly closed. Store at controlled room temperature 15° to 30° C (59° to 86° F). U.S. Patent No. 3,997,052 815 831 001 The Ujohn Co. Kalamazoo, MI 49001, USA


Halcion® *Ujohn Co.*
Tablets
triazolam
tablets, USP
0.125mg

Caution: Federal law prohibits dispensing without prescription.

NDC 0009-0010-37
10 Tablets

0009-0010-37 9

29. Mr. Bates has a history of seizure activity. Phenobarbital 15 mg po q 3 h is ordered. How many tablets will the nurse administer per dose? _____



NDC 0002-1032-04
1000 TABLETS No. 1545

Lilly **TV**

PHENOBARBITAL TABLETS, USP
30 mg


WARNING—May be habit forming.
CAUTION—Federal (U.S.A.) law prohibits dispensing without prescription.

Keep Tightly Closed
Store at Controlled Room Temperature
59° to 86°F (15° to 30°C)

WV D681 AMX
Eli Lilly & Co., Indianapolis, IN 46285, U.S.A.
Expiration Date/Control No.

Usual Adult Sedative Dose—15 to 30 mg
2 to 4 times a day.
Usual Adult Hypnotic Dose—100 to 200 mg
Dispense in a tight container.

30. Mrs. Ortega has chronic sinusitis. Her licensed prescriber orders amoxicillin 125 mg po q 8 h. How many milliliters will the nurse administer per dose? _____



N0047-2501-18

RECONSTITUTE w/105 mL WATER

Dispensing Directions - Prepare suspension at time of dispensing. Shake bottle to loosen powder or break up the cake that may have formed. Add a total of 105 mL water in 2 portions and shake well after each. This provides 150 mL of suspension. Each 5 mL contains amoxicillin trihydrate equivalent to 250 mg amoxicillin. Usual Dosage - Patients weighing less than 20 kg (44 lb): 20 mg / kg / day in divided doses every eight hours. Patients weighing 20 kg or more: 250 mg every eight hours. See package insert. Bottle contains amoxicillin trihydrate equivalent to 7.5 g amoxicillin. Store below 30°C (86°F). Protect from moisture. Manufactured for: WARNER CHILCOTT LABS Div of Warner-Lambert Co. 1990 Morris Plains, NJ 07950 USA By: Clonmel Chemicals Co. Ltd. Clonmel, Republic of Ireland 2501.0031

SHAKE WELL BEFORE USING
Keep bottles tightly closed. Any unused portion must be discarded after 14 days. Refrigeration is preferable but not required. Be sure to take each dose prescribed by your physician. Keep this and all drugs out of the reach of children. This bottle will contain 150 mL when prepared for dispensing as directed. Extra space in the bottle allows for shaking. 2501G232


Amoxicillin for Oral Suspension, USP
250mg per 5 mL
when reconstituted according to directions
Caution - Federal law prohibits dispensing without prescription.
150 mL bottle

WARNER CHILCOTT

TEAR ALONG HERE
6505-01-056-4195

Lot and Exp. date of powder

31. The licensed prescriber prescribes Tenormin 25 mg po q 4 h for Mr. Hutton's high blood pressure. How many tablets will you administer per dose? _____



NDC 0310-0105-10 100 tablets

TENORMIN®
(atenolol)

Rx only
Manufactured for:
AstraZeneca Pharmaceuticals LP
Wilmington, DE 19850
By: IPR Pharmaceuticals, Inc.
Carolina, PR 00984

USUAL DOSAGE: One tablet daily, or as directed by physician. Professional Information Brochure attached.
WARNING: As with all medications, keep out of the reach of children.
Store at controlled room temperature, 20-25°C (68-77°F) [see USP]. Dispense in well-closed, light-resistant containers. All trademarks are the property of the AstraZeneca group. 411308

LOT: _____
EXP: _____

AstraZeneca

F.P.O.
0310-0105-10 2

32. The licensed prescriber orders quinidine 0.6 g po q 4 h. Quinidine is supplied in 200-mg tablets. How many tablets will you give for one dose? _____ How many tablets will you give in 24 hours? _____

33. Mrs. Farmer has Zofran 8 mg po three times a day ordered for relief of nausea. How many milliliters will the nurse administer per dose? _____

NDC 0173-0489-00 *GlaxoWellcome*

Zofran®
(ondansetron hydrochloride)
Oral Solution

4 mg/5 mL

Caution: Federal law prohibits dispensing without prescription.


50 mL

Each 5 mL contains 5 mg of ondansetron HCl dihydrate equivalent to 4 mg of ondansetron.


See package insert for Dosage and Administration.

Store between 15° and 30°C (59° and 86°F). Protect from light. Store bottle upright in carton.

Glaxo Wellcome Inc.
Research Triangle Park, NC 27709
Made in England
4058968
Rev. 3/96



34. Your patient has Cipro 750 mg po q 12 h ordered for a severe respiratory tract infection. You have Cipro oral suspension 500 mg/5 mL available. How many milliliters will you administer per dose? _____



35. Mr. Golden, recovering from a left great toe amputation, has Colace elixir 100 mg po at bedtime ordered for constipation. How many milliliters will the nurse administer per dose? _____

NDC 0087-0720-01

Do not use if carton overwrap was missing or broken.
COLACE is used for prevention of dry, hard stools.
Usual daily dose
Infants and children under 3: As prescribed by physician.
Children 3 to 6: 1 to 3 teaspoons.
Children 6 to 12: 2 teaspoons one to three times daily.
Adults and older children: 1 to 3 tablespoons.
Keep this and all medication out of reach of children.

SYRUP
COLACE®
DOCUSATE SODIUM
STOOL SOFTENER

8 FL OZ (1/2 PT)

Mead Johnson


The effect of COLACE on the stools may not be apparent until 1 to 3 days after first oral dose.

WARNING: As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.

Each teaspoon (5 ml) contains 20 mg docusate sodium; each tablespoon (15 ml) contains 60 mg. Contains not more than 1% alcohol.

Store at room temperature. Protect from excessive heat.

MEAD JOHNSON
PHARMACEUTICALS
Bristol-Myers
U.S. Pharmaceutical and Nutritional Group
Evansville, IN 47721
Made in U.S.A.



36. Mr. Malito is to receive Procanbid 1 g po now. How many tablets will you administer per dose? _____

Exp. date and lot
0564G110

Each tablet contains 500 mg of procainamide hydrochloride. Usual Dosage—See package insert for complete prescribing information. Keep this and all drugs out of the reach of children. Store at controlled room temperature, 20°–25° C (68°–77° F). Do not store above 30° C (86° F). Procainamide is not light-resistant. See USP *Controlled Room Temperature*.

NDC 61570-069-60

Procanbid®
[Procainamide HCl
Extended-Release Tablets*]

500 mg

Rx Only
60 TABLETS

**Monarch
Pharmaceuticals®**

Distributed by: Monarch Pharmaceuticals, Inc., Bloomington, TN 37190
Manufactured by: Warner-Lambert Co., Morris Plains, NJ 07950

3 61570 06960 17

37. Mr. Mikal was admitted for treatment of leukemia and receives Deltasone 7.5 mg po three times a day as part of his chemotherapy. The drug is available in 2.5-mg tablets. How many tablets will the nurse administer per dose? _____

38. The licensed prescriber orders Tegretol 0.2 g po three times a day for Mr. Pine's epilepsy. How many tablets will the nurse administer per dose? _____

EXP LOT

Do not store above 89°F (30°C).
Dosage: See package insert.
Protect from light and moisture.
Ciba-Geigy Corporation
Pharmaceuticals Division
Summit, NJ 07901

NDC 0083-0052-30 FSC 1821
6505-01-153-4524

Tegretol® 100 mg
carbamazepine USP

Chewable Tablets

100 tablets
Dispense in tight, light-resistant container (USP).
Caution: Federal law prohibits dispensing without prescription.

Keep this and all drugs out of the reach of children.

3 0083-0052-30 0

643330

39. Lortab 5/500 po now is ordered for Mrs. Lindl for pain. How many tablets will she receive per dose? _____ How much acetaminophen is in each tablet? _____

ucb Pharma

PHARMACIST: Dispense in a tight, light-resistant container with a child-resistant closure.
Store at controlled room temperature, 15°–30° C (59°–86°F).

Lot No.:
Exp. Date:

Manufactured for
UCB Pharma, Inc.
Synovis, CA 30090
by Mallinckrodt Inc.
Hobart, NY 13788

NDC 50474-902-01 100 TABLETS

LORTAB 5/500

**HYDROCODONE BITARTRATE
AND ACETAMINOPHEN
TABLETS, USP**
5 mg/500 mg

Each scored, white with blue specks tablet contains:
Hydrocodone Bitartrate 5 mg
Acetaminophen 500 mg

Rx only


USUAL DOSAGE: See package insert for complete dosage recommendations.

3 50474-902-01 1

Rev 6/01
P/N 1003723

H


40. Mrs. Cross was admitted with a myasthenia crisis. Decadron 0.5 mg po q 12 h is ordered. How many tablets will the nurse administer per dose? _____



41. Mr. Cook requires medication for nausea. Compazine 10 mg po q 4 h prn is ordered. You have Compazine 5-mg tablets available. How many tablets will you administer per dose? _____

42. Mr. Pace receives Atarax 100 mg po at bedtime prn to relieve anxiety. You have 50-mg tablets available. How many tablets will you administer per dose? _____

43. The licensed prescriber orders Rifadin 600 mg po 1 hour before dinner daily. How many capsules will the nurse administer per dose? _____



44. Mr. Day receives Lanoxin 0.25 mg po daily for atrial fibrillation. How many tablets will the nurse administer per dose? _____



45. Mr. Payne receives Keflex 500 mg po four times a day before his dental extraction. How many capsules will the nurse administer in each dose? _____ How many capsules will the nurse administer for 1 day? _____

NDC 0777-0869-02
100 PULVULES[®] No. 402

KEFLEX[®]
CEPHALEXIN
CAPSULES, USP
250 mg

CAUTION—Federal (U.S.A.) law prohibits dispensing without prescription. Keep Tightly Closed. Store at Controlled Room Temperature 59° to 86°F (15° to 30°C). WV 3473 DPK. Manufactured by DISTA PRODUCTS CO., a Division of Eli Lilly and Company, Indianapolis, IN, U.S.A. Expiration Date/Control No.

Usual Adult Dose—One PULVULE every 6 hours. For more severe infections, dose may be increased, not to exceed 4 g a day. See literature. Each PULVULE contains Cephalixin Monohydrate equivalent to 250 mg Cephalixin. Dispense in a tight container.

3 0777 0869 02 9

46. Mr. Tune is experiencing gastroesophageal reflux. The licensed prescriber orders Nexium 40 mg po daily for 3 days. How many capsules will the nurse administer per dose? _____

NDC 0186-5020-54

Nexium[™]
(esomeprazole magnesium)
90 Delayed-Release Capsules
Rx only
*Each delayed-release capsule contains 20 mg esomeprazole

Keep container tightly closed. Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F). (See USP Controlled Room Temperature). Dispense in a tight container. USUAL ADULT DOSAGE: See package insert. All trademarks are the property of the AstraZeneca group. © AstraZeneca 2000. Manufactured for: AstraZeneca LP, Wilmington, DE 19850. By: AstraZeneca AB, S-151 85 Södertälje, Sweden. Product of France.

20 mg

AstraZeneca

9347701
90 | No. 5020 620309-03

47. Mrs. Graves receives phenobarbital tablets 90 mg po q 3 h prn for seizure activity. How many tablets will the nurse administer per dose? _____

NDC 0002-1032-04
1000 TABLETS No. 1545

Lilly **IV**

PHENOBARBITAL
TABLETS, USP
30 mg

WARNING—May be habit forming.

CAUTION—Federal (U.S.A.) law prohibits dispensing without prescription.

Usual Adult Sedative Dose—15 to 30 mg 2 to 4 times a day.
Usual Adult Hypnotic Dose—100 to 200 mg
Dispense in a tight container.

Keep Tightly Closed. Store at Controlled Room Temperature 59° to 86°F (15° to 30°C). WV 0681 ANX. Eli Lilly & Co., Indianapolis, IN 46285, U.S.A. Expiration Date/Control No.

48. Mr. Vee is enrolled in a smoking-cessation program. He is to begin with Wellbutrin 150 mg po daily for 3 days. How many tablets will the nurse administer per dose? _____

100 Tablets NDC 0173-0177-55 NDC 0173-0177-55
 Each film-coated tablet contains
75 mg
WELLBUTRIN[®]
 (bupropion hydrochloride)
Tablets

WARNING: Do not use in combination with ZYBAN[®], or any other medicines that contain bupropion hydrochloride.
 Store at 15° to 25°C (59° to 77°F). Protect from moisture.
 Dispense in a tight, light-resistant container as defined in the USP
 Rx only

GlaxoWellcome

Manufactured by
 Catalytic Pharmaceutical, Inc
 Greenville, NC 27834
 for Glaxo Wellcome Inc
 Research Triangle Park, NC 27709

4117573 Rev. 9/89

NSN 6505-01-314-6641

3 0173-0000-00 2

646124

LOT
 EXP

49. Mr. Sahl, recovering from a coronary artery bypass graft, receives Capoten 2.5 mg po twice a day. How many tablets will the nurse administer per dose? _____

Exp.:
 Cont.:
 Keep tightly closed (protect from moisture)
 Do not store above 86°F NDC 0003-0450-54

E. R. Squibb & Sons, Inc.
 Princeton, NJ 08543 USA
 Made in USA 5131600R-1

SQUIBB

100 tablets NDC 0003-0450-54

12.5 mg
CAPOTEN[®]
 Captopril Tablets

Caution: Federal law prohibits dispensing without prescription
 6505-01-226-2924

Each tablet contains 12.5 mg captopril
 Dosage: See insert
 US Patent: 4,105,776

TO PEEL—LIFT HERE →

50. Mr. Dale receives Zantac 150 mg two times a day as part of his treatment for esophagitis. How many milliliters will the nurse administer per dose? _____

Glaxo Pharmaceuticals

Zantac
 (ranitidine hydrochloride)
 Syrup
 15 mg/mL

Caution: Federal law prohibits dispensing without prescription.

16 fl oz (1 pint)

LOT
 EXP